

5

## Referências bibliográficas

ABDULRAHMAN A. A., SAAD A. A., AKEL A. A. Flow-injection determination of kanamycin by inhibition of the lucigenin-H<sub>2</sub>O<sub>2</sub>-Co<sup>2+</sup> system **Anal. Chim. Acta.** V. 292, p. 201-208, 1994.

AGUI, M. L., GARCÍA-CAMPANA, A. M., GÁMIZ-GRACIA, L., BLANCO, C. C. Laser induced fluorescence coupled to capillary electrophoresis for the determination of fluoroquinolones in foods of animal origin using molecularly imprinted polymers. **J. Chromatogr. A.** V. 1217, p. 2237–2242, 2010.

ALI, W. H., D. DELPHINE, ALIX, F., C. PÉROLIER, O. LÉPINE, S. BAYOUDH,. Solid-phase extraction using molecularly imprinted polymers for selective extraction of a mycotoxin in cereals. **J. Chromatogr. A.** V. 1217, p. 6668–6673, 2010.

ALKINDY, S.; BADIA, R.; SUAREZ-RODRIGUEZ, J. L.; D., M. E.; **Crit. Rev. Anal. Chem.** V. 30, 291, 2000.

ALVES, A.M.P.; ALVES, E.P.B.; BUTTOW, P., J.V.C.M; ZANONI, J.N.; STABILE, S.R.; ANDERSON, L.I.; B. SELLERGEN; K. MOSBACH. Imprinting of amino acid derivatives in macroporous polymers, **Tetrahedron Lett.** V.25, p. 5211, 1984.

ANVISA. Agência Nacional de Vigilância Sanitária. **Resolução RE nº 899 de 29 de maio de 2003 a.** Guia para a Validação de Métodos Analíticos e Bioanalíticos.

ARIFFIN, M.M.; E. I. MILLER; P. A. G. COMACK; R. A. ANDERSON. Molecularly Imprinted Solid-Phase Extraction of Diazepam and Its Metabolites from Hair Samples, **Anal. Chem.** V. 79 p. 256, 2007.

Aspectos gerais e abordagem terapêutica da quercetina sobre as complicações do diabetes causadas pelo estresse oxidativo. **Arq. Ciênc. Saúde Unipar.** V.14, n.2, p. 179-186, 2010.

BALAMURUGAN, K., KANNAN G., TANGIRALA P. Preparation and evaluation of molecularly imprinted polymer liquid chromatography column for the separation of Cathine enantiomers. **Saudi Pharmaceut. J.** V.20, p1. 53-61, 2012.

BEHLING, E.B., SENDÃO, M. C., FRANCESCATO, H. D. C., ANTUNES, L. M. G., BIANCHI, M. L. Flavonóide Quercetina: Aspectos gerais e ações biológicas. **Alim. Nutr.** V. 15, n. 3, p. 285-292, 2004.

BEVILÁQUA, R. Resumo de farmacologia aminoglicosídios. Penildon – capítulo 104.

BRAVO,J.C.; R.M. GARCINUNO; P. FERNANDEZ; J. S. DURANO. A new molecularly imprinted polymer for the on-column solid-phase extraction of diethylstilbestrol from aqueous samples, **Anal. Bioanal. Chem.** V.388 p.1039, 2007.

BUENO, M.J. MARTÍNEZ, S. HERRERA, A. UCLÉS, A. AGUERA, M. D. HERMANDO, O. SHIMELIS, M. RUDOLFSSON, A. R. FERNANDEZ-ALBA. Determination of malachite green residues in fish using molecularly imprinted solid-phase extraction followed by liquid chromatography–linear ion trap mass spectrometry. **Anal. Chim. Acta.** V.665, p. 47–54, 2010.

BUTTON, S. TONINI. Metodologia para planejamento experimental e análise de resultados. Universidade Estadual de Campinas, 2012.

CABRAL, A.P.G. de A. **Estudo de polímeros de impressão molecular e sua aplicação na determinação da morfina.** Universidade Estácio de Sá, 2008. (Monografia de final de curso).

CÉSAR-OLIVEIRA, M. A. F., S. ZAIONCZ, A. R. S. OLIVEIRA, M. C. R. ALMEIDA, S. F. ZAWADZKI, L. AKCELRUD, M. AGUIAR, D. TABAK e E. F. LUCAS. Síntese de Copolímeros Metacrílicos Através da Modificação Química do

Poli (Metacrilato de Metila) de Massa Molar Controlada. **Polímeros: Ciência e Tecnologia – Artigo técnico científico.** Out/Dez – 1999.

CHEN, L., X. ZHANG, Y. XU,. Determination of fluoroquinolone antibiotics in environmental water samples based on magnetic molecularly imprinted polymer extraction followed by liquid chromatography–tandem mass spectrometry. **Anal. Chim. Acta.** V. 662, p. 31–38, 2010.

CHEN, X., Z.. ZHANG, X.Y., J. LI , Y. LU , H. j. CHEN , W. RAO, S. YAO. Molecularly imprinted polymers based on multi-walled carbon nanotubes for selective solid-phase extraction of oleanolic acid from the roots of kiwi fruit samples. **Talanta.** V. 99, p. 959–965, 2012.

COMPLETO, C.D. da SILVA. Adsorção de albumina bovina e ácido hialurônico em cerâmicos de titânio. Universidade Técnica de Lisboa, 2008, (Dissertação de Mestrado).

DE BOER, V. C. Tissue distribution of quercetin in rats and pigs. **J. Nutr.** v. 135, n. 7, p. 1718-1725, 2005.

DENG, Q.L, Y. LI, L. HUA Z., Y. K. ZHANG. Molecularly imprinted macroporous monolithic materials for protein recognition. **Chin. Chem. Lett..** V. 22, p. 1351–1354, 2011.

DIAS, A. S. O antioxidante queracetina diminui o estresse oxidativo hepático em ratos diabéticos. Universidade Federal do Rio Grande do Sul. 2009. Tese de Doutorado.

DICKEY, F.H. The Preparation of specific adsorbents. **Proc. Natl. Acad. Sci.,** v. 35, n. 5, p. 227-229, 1949.

DUAN, J. L. S., J. ZHA. One-Pot Synthesis of Highly Luminescent CdTe Quantum Dots by Microwave Irradiation Reduction and Their  $Hg^{2+}$ -Sensitive Properties.

DUFOR C., O. DANGLES. Flavonoid–serum albumin complexation: determination of binding constants and binding sites by fluorescence spectroscopy. **Biochim. Biophys. Acta.** V. 1721 p. 164– 173, 2005.

FERNANDEZ-LIANO, L., M.C. B.Lopez, M.J. L. CASTANON, A.J. M. ORDIERES, P. T.BLANCO. Determination of diclofenac in urine samples by molecularly-imprinted solid-phase extraction and adsorptive differential pulse voltammetry, **Electroanalysis**. V. 19 p.1555, 2007.

FERREIRA, E. S. Interação da albumina do soro bovino (BSA) com substratos sintéticos. Universidade de São Paulo, 2009, (Tese de Doutorado).

FIGUEIREDO, C. de, E. Emprego de polímeros de impressão molecular (MIP) na extração e pré-concentração de analitos orgânicos em amostras biológicas seguido de determinação espectrofotométrica. Universidade Estadual de Campinas,2009, (Tese de Doutorado).

FIGUEIREDO, E. C., A. C. B. DIAS, M. A., Z. ARRUDA. Impressão molecular: uma estratégia promissora na elaboração de matrizes para a liberação controlada de fármacos. **Revista Brasileira de Ciências Farmacêuticas Brazilian Journal of Pharmaceutical Sciences**.V. 44, jul./set., 2008.

FILHO, A.M.; F. N. SANTOS; P. O. A. de P. PEREIRA. Development, validation and application of a methodology based on solid-phase micro extraction followed by gas chromatography coupled to mass spectrometry (SPME/GC–MS) for the determination of pesticide residues in mangoes. **Talanta**. V. 81, p. 346–354, 2010.

FRANCES, A., E.TURRILLAS; J. V. MERCADER, C. AGULLÓ; A. ABAD-SOMOVILLA, A. ABAD-FUENTES. Development of immunoaffinity columns for pyraclostrobin extraction from fruit juices and analysis by liquid chromatography with UV detection. **J. Chromatogr. A**. V. 1218, p. 4902– 4909, 2011.

GONZALES-RODRIGUEZ, R.M.; BEATRIZ CANCHO-GRANDE; J. S.GANDARA. Multiresidue determination of 11 new fungicides in grapes and wines by liquid-liquid extraction/clean-up and programmable temperature vaporization injection with analyte protectants/gas chromatography/ion trap mass spectrometry. **J. Chromatogr. A**. V. 1216, p. 6033–6042, 2009.

GRASSI, V.. Polímeros molecularmente impressos (MIPs) como extratores em fase sólida em sistemas de análises em fluxo. Universidade de São Paulo. 2008. (Tese de Doutorado).

HAGINAKA, J., C. KAGAWA. Retentivity and enantioselectivity of uniformly sized molecularly imprinted polymers for d-chlorpheniramine and -brompheniramine in hydro-organic mobile phases, **J. Chromatogr. B.** v. 804 p.19, 2004.

HAVSTEEN, B.N. The biochemistry and medical significance of the flavonoids. **Pharmacol. Therapeut.** V. 96, p. 67-202, 2002.

HOSHINA, K., S. HORIYAMA, H. MATSUNAGA, J. HAGINAKA. Simultaneous determination of non-steroidal anti-inflammatory drugs in river water samples by liquid chromatography–tandem mass spectrometry using molecularly imprinted polymers as a pretreatment column. **J. Pharmaceut. Biomed.** V. 55, p. 916–922, 2011.

HOSOYA, K.; Y., K.; TANAKA, N.; KIMATA, K.; ARAKI, T.; HAGINAKA. **J. Chem. Lett.**,v.8, p. 1437, 1994.

HUANG, B., Y. CHEN, G. WANG, C. LIU. Preparation and evaluation of a monolithic molecularly imprinted polymer for the chiral separation of neurotransmitters and their analogues by capillary electrochromatography. **J. Chromatogr A.** V. 1218, p. 849–855, 2011.

HUBER, L. SENGER, D. B. RODRIGUEZ-AMAYA. Flavonóis e flavonas: fontes brasileiras e fatores que influenciam a composição em alimentos. **Alim. Nutr.** V. 19, p.97-108 , 2008.

HWANG, C., W. L., Chromatographic resolution of the enantiomers of phenylpropanolamine by using molecularly imprinted polymer as the stationary phase, **J. Chromatogr. B.** V. 765 p. 45, 2001.

INMETRO. Instituto Nacional de Metrologia, Qualidade e Tecnologia. **Orientação sobre validação de métodos analíticos.** DOQ-CGCRE-008 Revisão 04 – jul/2011 a.

INMETRO. Instituto Nacional de Metrologia, Qualidade e Tecnologia. **Orientação sobre validação de métodos analíticos.** DOQ-CGCRC-008 Revisão 04 – jul/2011 a.

JAVANBAKTA, M. A. M. ATTARAN, M. H. NAMJUMANESH, M. E. MANESH, B. A. ADERGANI. Solid-phase extraction of tramadol from plasma and urine samples using a novel water-compatible molecularly imprinted polymer. **J. Chromatogr B.** V. 878, p. 1700–1706, 2010.

JAYABHARATHI, J., VENUGOPALI T.; MARIMUTHU V.; KANAGARATHINAM S. Displacement Reaction Using Ibuprofen in a Mixture of Bioactive Imidazole Derivative and Bovine Serum Albumin—a Fluorescence Quenching Study. **J. Fluoresc.** V. 21, p. 1825-1830, 2011.

JHONSI, M.A.; ARUNKUMAR K., RAJALINGAM R., MAHMMOUD, S. A E. SADEK. Photoinduced Interaction of MPA-Capped CdTe Quantum Dots with Denatured Bovine Serum Albumin. **Nanosc. and Nanotechnol. Lett..** V. 3, p.125–130, 2011.

JOKLIK, W. K.; PHIL, D. Virology. cap. 22- Arboviruses. 2. ed. Norwalk: AppletonCentury-Crofts, 1985.

JULIATTI, F. C. Modo de ação dos fungicidas sobre plantas e fungos. Departamento de fitopatologia, ICIAG/Universidade Federal de Uberlândia. Resumo, 2005.

KAALE, E.; A. VAN S.; E. ROETS, J. H. Development and validation of a simple capillary zone electrophoresis method for the analysis of kanamycin sulfate with UV detection after pre-capillary derivatization. **J. Chromatogr A.** V. 924, p. 451–458, 2001.

KALDAS M. I.; U. K. WALLE; H. V. D. WOUDE; J. M. MCMILLAN; T. WALLE. Covalent Binding of the Flavonoid Quercetin to Human Serum Albumin. **J. Agric. Food Chem.**, v.53, p.4194-4197, 2005.

KAN, X., Z. XING, A. ZHU, Z. ZHAO, G. XU, CHEN L., H. ZHOU. Molecularly imprinted polymers based electrochemical sensor for bovine hemoglobin recognition. **Sensor Actuator B.** V. 168, p.395-401, 2012.

KEMPE, M.; **Anal. Chem.**, v.68, p.1948, 1996.

KIM, B, HYOUN, Y. KIM, J. H. OK. Development of liquid chromatographic method for the analysis of kanamycin residues in varicella vaccine using phenylisocyanate as a derivatization reagent. **J. Chromatogr B.** V.752, p. 173–177, 2001.

KIM, H., G. G. Comparison of the Thermodynamic Properties of Particulate and Monolithic Columns of Molecularly Imprinted Copolymers, **Anal. Chem.** V. 77 p.93, 2005.

KITSON, T. M. Spectrophotometric and kinetic studies on the binding of the bioflavonoid quercetin to bovine sérum albumin. **Biosci. Biotechnol. Biochem.** V. 68, n°10, p. 2165-2170, 2004.

KUMAR, A.; A. K. M., DHANANJAY K. T. A new method for determination of myricetin and quercetin using solid phase microextraction–high performance liquid chromatography–ultra violet/visible system in grapes, vegetables and red wine samples **Anal. Chim. Acta.** V. 631, p. 177–181, 2009.

KUMAR, D., BHIM B. Prasad. Multiwalled carbon nanotubes embedded molecularly imprinted polymer-modified screen printed carbon electrode for the quantitative analysis of C-reactive protein. **Sensor and Actuator B.** V. 171– 172, p. 1141– 1150, 2012.

LAGUNAS-ALLUE, L., M.T. M., JESUS S.; ARNAUD S.; CORINE F. JEAN M. C. Degradation intermediates and reaction pathway of pyraclostrobin with TiO<sub>2</sub> photocatalysis. **Applied Catalysis B: Environmental.** V. 115-116, p. 285– 293, 2012.

LAKOWICZ, J. Principles of fluorescence Spectroscopy, third ed., Springer, New York, 2006.

LEE, T. P., B. SAAD, W. S. KHAYOON, B. SALLEH. Molecularly imprinted polymer as sorbent in micro-solid phase extraction of ochratoxin A in coffee, grape juice and urine. **Talanta**. V. 88, p. 129– 135, 2012.

Levy, J. A.; Fraenkel-conrat, H.; Owens, R. A. *Virology*. 3 ed. New Jersey: Prentice Hall, 1994.

LIANDA, R.L.P. Perfil de substâncias fenólicas de méis brasileiros por cromatografia líquida de alta eficiência e avaliação do potencial antioxidante. Universidade Federal Rural do Rio de Janeiro, 2009, (Tese de Doutorado).

LIANG L.; HUIFENG Q.; NENGHU U F.; JICUN R. Significant enhancement of the quantum yield of CdTe nanocrystals synthesized in aqueous phase by controlling the pH and concentrations of precursor solutions. **J. Luminesc.** V.116, p. 59–66, 2006.

LIBARDI, S. H.. Atividade antioxidante da vanilina e do ácido vanílico e o efeito da complexação por proteínas do soro do leite na desativação de radicais e ferrilmiooglobina em condições simulando o trato gastrointestinal. Universidade de São Paulo, 2010, (Dissertação de Mestrado).

LIN, J.; T. N., K. UCIYAMA; T. HOBO. Molecularly imprinted polymer as chiral selector for enantioseparation of amino acids by capillary gel electrophoresis Chromatographia. V. 43 p. 585, 1996.

LIU, E-Hu; L. QI; P. LI. Structural Relationship and Binding Mechanisms of Five Flavonoids with Bovine Serum Albumin. **Molecules**. V. 15, p. 9092-9103, 2010.

LIU, X., Z. Y.HUANG, J. YANG, Z.S. LIU. Molecularly imprinted nanoparticles with nontailing peaks in capillary Electrochromatography. **J. Chromatogr. A**. V. 1264, p. 137– 142, 2012.

LONG, Y.H.; M. H.ERNANDEZ, E. KAALE; A. V. SCHEPDAEL, E. ROETS, F. BORRULL, M. C.L; J. HOOGMARTENS. Determination of kanamycin in serum by solid-phase extraction, pre-capillary derivatization and capillary electrophoresis. **J. Chromatogr B**. V.784, p. 255–264, 2003.

LÓPEZ, M.D. M. C., M.C. C. PÉREZ, M. S. D. GARCÍA, J. M. L. VILARINO, M. V. G. RODRIGUEZ, L. F. B. LOSADA. Preparation, evaluation and characterization of quercetin-molecularly imprinted polymer for preconcentration and clean-up of catechins. **Anal. Chim. Acta.** V. 721 p.68 – 78, 2012.

LUCCI, P., D. DERRIEN, F. ALIX, C. PÉROLIER, S. BAYOUDH. Molecularly imprinted polymer solid-phase extraction for detection of zearalenone in cereal sample extracts. **Anal. Chim. Acta.** V. 672, p. 15-19, 2010.

LUO, X., Y. ZHA, X. TU, Y.HUAN, S. LUO, L. YAN. Novel molecularly imprinted polymer using 1-(alfa-methyl acrylate)-3-methylimidazolium bromide as functional monomer for simultaneous extraction and determination of water-soluble acid dyes in wastewater and soft drink by solid phase extraction and high performance liquid chromatography. **J. Chromatogr. A.** v. 1218, p. 1115–1121, 2011.

MACHADO, F., L., E. L., PINTO, J. C. Uma Revisão Sobre os Processos de Polimerização em Suspensão. **Polímeros: Ciência e Tecnologia.** V. 17, p. 166-179, 2007.

MAHONY, J.O.; A. MOLINELLII B.; K. NOLAN A.; M.R. SMYTH; B. MIZAIKOFF. Anatomy of a successful imprint: Analysing the recognition mechanisms of a molecularly imprinted polymer for quercetin. **Biosens. Bioelectron..** V. 21 p.1383–1392. 2006.

MAHONY, J.O.; WEI S.; Molinelli, A.; MIZAIKOFF, B. Imprinted Polymeric Materials. Insight into the Nature of Prepolymerization Complexes of Quercetin Imprinted Polymers **Anal. Chem.** V. 78, p. 6187-6190, 2006.

MARTÍN-ESTEBAM, A. Molecularly imprinted polymers: new molecular recognition materials for selective solid-phase extraction of organic compounds, **Fresenius J. Anal. Chem.** V.370 p. 795, 2001.

MATSUBARA, S., D. B. RODRIGUEZ-AMAYA. Conteúdo de mirocetina, quercetina e kampferol em chás comercializados no Brasil. **Ciênc. Tecnol. Aliment.** V. 26, p. 380-385, 2006.

MATSUI, J., OKADA, M. TSURUOKA, M., TAKEUCHI, T. **Anal. Commun.**, v.34, p.85, 1997.

MAYES, A.G.; L. I. AANDERSON, K. MOSBACH. Sugar binding polymers showing high anomeric and epimeric discrimination obtained by noncovalent molecular imprinting, **Anal. Biochem.** V. 222 p. 483, 1994.

MAYS, D. L., RICHARD J. V. APELDOORN, RONALD G. LATIACK. High-performance liquid chromatographic determination of kanamycin. **J. Chromatogr.** V. 120, p. 93-102, 1976.

MEGOULAS, N.C., KOUPPARIS M.A. Direct determination of kanamycin in raw materials, veterinary formulation and culture media using a novel liquid chromatography–evaporative light scattering method. **Anal. Chim. Acta**. V. 547, p. 64–72, 2005.

MHAKA, B., E. CUKROWSKA, B. T. S. BUI, O. RAMSTROM, K. HAUPT, H. TUTU, L. CHIMUKA. Selective extraction of triazine herbicides from food samples based on a combination of a liquid membrane and molecularly imprinted polymers. **J. Chromatog. A**. v. 1216, p.6796–6801, 2009.

MISHA, B., A. BARIK, K. I. PRIYADARSINI, H. MOHAN. Fluorescence spectroscopic studies on binding of a flavonoid antioxidant quercetin to serum albumins. **J. Chem. Sci.** V. 117, Nº. 6, p. 641–647, 2005.

MOEIN, M. MAHDI, M. JAVANBAKHT, B. A.. Molecularly imprinted polymer cartridges coupled on-line with high performance liquid chromatography for simple and rapid analysis of dextromethorphan in human plasma samples. **J. Chromatogr. B**. V. 879, p.777–782, 2011.

MOLINA, C. Q., B. CLAUDE, A. M. G. M. del O. , P. MORIN. Convenient solid phase extraction of cephalosporins in milk using a molecularly imprinted polymer. **Food Chemistry**. V. 135, p. 775–779, 2012.

MOLINELLI, A.; WEISS, R. Mizaikoff, B. Advanced Solid Phase Extraction Using Molecularly Imprinted Polymers for the Determination of Quercetin in Red Wine.,**J. Agric. Food Chem.** V. 50, p. 1804-1808, 2002.

Muhammad,T., L. CUI, W. JIDE, E. V. PILETSK, A. R. GUERREIRO, S. A. PILETSKY. Rational design and synthesis of water-compatible molecularly imprinted polymers for selective solid phase extraction of amiodarone. **Anal. Chim. Acta.** V.709, p. 98– 104, 2012.

NAVARRO-VILLOSLAD, F. **Anal. Chim. Acta.** V.504, p.149, 2004.

NETO, J. de R. M. Sensor amperométrico a base de um polímero de impressão molecular com protoporfirina IX de ferro para a determinação de 4-aminofenol. Universidade Federal do Maranhão. 2010. Tese de Mestrado.

NIELSEN, S.E. L.O. DRAGSTED. Column-switching high-performance liquid chromatographic assay for the determination of quercetin in human urine with ultraviolet absorbance detection, **J. Chromatogr. B.** V.707 p. 81–89, 1998.

OLIVEIRA, C. A.. Métodos analíticos baseados na luminescência para a determinação das fluorquinolonas esparfloxacina e enoxacina. Pontifícia Universidade Católica – PUC-Rio, 2010 (Dissertação de Mestrado).

OLIVEIRA, J.F.P., J. P. CIPULLO, E. A. BURDMANN. Nefrotoxicidade dos aminoglicosídeos. **Braz J Cardiovasc. Surg.** V. 21, p. 444-452, 2006.

OU, J.J.; X. LI; S. FENG, J. DONG, X. DONG, L. KONG; M. YE; H. ZOU. Preparation and evaluation of a molecularly imprinted polymer derivatized silica monolithic column for capillary electrochromatography and capillary liquid chromatography, **Anal. Chem.** V. 79 p.639, 2007.

OU, J., H. LINA, S. TANG, Z. ZHANGA, J. DONG, H. ZOU. Hybrid monolithic columns coated with cellulose tris(3,5-dimethylphenyl-carbamate) for enantioseparations in capillary electrochromatography and capillary liquid chromatography. **J. Chromatogr. A.** V. 1269, p.372– 378, 2012.

PAKADE, V., S. LINDAHL, L. CHIMUKA, C. TURNER. Molecularly imprinted polymers targeting quercetin in high-temperature aqueous solutions. **J. Chromatogr. A.** V. 1230 p. 15– 23, 2012.

PAPAPPOULOU, A., R J. GREEN, R. A. FRAZIER. Interaction of Flavonoids with Bovine Serum Albumin: A Fluorescence Quenching Study. **J. Agric. Food Chem.** V. 53, p. 158-163, 2005.

PAULLING, L.J. A Theory of the Structure and Process of Formation of Antibodies. **J. Am. Chem. Soc.** V. 62, n. 783, p. 2643-2657, 1940.

PENILDON, S. Farmacologia. 6 ed. Rio de Janeiro: Guanabara Koogan, 2002.

PEREIRA, L. A. Desenvolvimento de um Método de extração em fase sólida molecularmente impressa (MISPE) para a determinação de fenantrotiona em tomate. Universidade Estadual de Campinas. 2008. Dissertação de Mestrado.

PLUNKETT, S., ARNOLD, F. **J. Chromatogr. A**, v.708, p.19, 1995.

PRASAD, B.B.; P. S. SHARMA, D. LAKSHMI. Molecularly imprinted polymer-based solid-phase extraction combined with molecularly imprinted polymer-based sensor for detection of uric acid, **J. Chromatogr. A**. V. 1173 p. 18, 2007.

QU, P., J. LEI, L. ZHANG, R. OUYANG, H. JU. Molecularly imprinted magnetic nanoparticles as tunable stationary phase located in microfluidic channel for enantioseparation. **J. Chromatogr. A**. V. 1217, p. 6115–6121, 2010.

QUAGLIA, M., D.E LORENZI, E., SULITZKY, C. MASSOLINI G. SELLERGEN, B. **Analyst**, v. 126, p.1495, 2001.

RAMSTROM, O., I. A. NICHOLLS, K. MOSBACH. Synthetic peptide receptor mimics -highly stereoselective recognition in noncovalent molecularly imprinted polymers, **Tetrahedron Asymmetr.** V. 5 p. 649, 1994.

RAO, R. N., P. K. MAURYA, S. KHALID. Development of a molecularly imprinted polymer for selective extraction followed by liquid chromatographic determination of sitagliptin in rat plasma and urine. **Talanta**. V. 85, p. 950– 957, 2011.

RIBANI, R. H., D. B. RODRIGUEZ-AMAYA. Otimização de método para determinação de flavonóis e flavonas em frutas por cromatografia líquida

de alta eficiência utilizando delineamento estatístico e análise de superfície de resposta. **Quím. Nova.** V.31, 1378-1384, 2008.

RODRIGUES, M. A. T. Avaliação do efeito fisiológico do uso de fungicidas na cultura de soja. Universidade de São Paulo, 2009, (Tese de Doutorado).

ROY, A. S., D. R. TRIPATHY, A. CHATTERJEE, S. DASGUPTA. A spectroscopic study of the interaction of the antioxidant naringin with bovine serum albumin. **J. Biophys. Chem.** Vol.1, N°. 3, p. 141-152, 2010.

RUSSO, F.M., S. C. TEDESCO, I., BIOTTO, S., R. G.L. The flavonoid quercetin in disease prevention and therapy: Facts and fancies **Biochem. Pharmac.** V. 83 p.6-15, 2012.

SANTOS, W. de J. R. Nanoreatores biomiméticos à peroxidase baseados em MIP: uma estratégia promissora para determinação de compostos fenólicos. Universidade Estadual de Campinas. 2009. Tese de Doutorado.

SCHWEITZ, L., L. I. ANDERSON, S. NILSSON. Capillary electrochromatography with predetermined selectivity obtained through molecular imprinting, **Anal. Chem.** V. 69 p.1179, 1997.

SCORRANO, S., L. LONGO, G. VASAPOLLO. Molecularly imprinted polymers for solid-phase extraction of 1-methyladenosine from human urine. **Anal. Chim. Acta.** V. 659, p. 167–171, 2010.

SELLERGEN, B. **Macromol. Chem. Phys.** v.190, p. 2703, 1989.

SELLERGEN, B. **Trends Anal. Chem.** V. 16, p.310, 1997.

SHE, Y., W. CAOA, X., SHIA, X. LV, J. LIU, R. WANGA, F. JIN , J. WANGA, H. XIAO. Class-specific molecularly imprinted polymers for the selective extraction and determination of sulfonylurea herbicides in maize samples by high-performance liquid chromatography–tandem mass spectrometry. **J. Chromatogr. B.** V. 878, p. 2047–2053, 2010.

SILVA, R. G. da C. Materiais sorventes impressos molecularmente preparados por processos sol-gel. Universidade Estadual de Campinas. 2009. (Tese de Doutorado).

SOLEIMANI, M., S. GHADERI, M. G. AFSHAR, S. SOLEIMANI. Synthesis of molecularly imprinted polymer as a sorbent for solid phase extraction of bovine albumin from whey, milk, urine and serum. **Microchem. Journal.** V. 100, p. 1–7, 2012.

SONG, W., J., ZHU, J. Effect of Porogenic Solvent on Selective Performance of Molecularly Imprinted Polymer for Quercetin. **Materials Research.** V. 12, N. 3, p.299-304, 2009.

SONG, X., J. LI, J. WANGA, LINGXIN CHEN. Quercetin molecularly imprinted polymers: Preparation, recognition characteristics and properties as sorbent for solid-phase extraction. **Talanta.** V. 80 p.694–702, 2009.

SOUZA, C. F de.; A. L. M. C. da CUNHA, R. Q. AUCÉLIO. Determination of Picoxystrobin and Pyraclostrobin by MEKC with On-Line Analyte Concentration. **Chromatogr.** V. 70, p. 1461–1466, 2009.

SOUZA, C. F. de. Desenvolvimento e aplicação de métodos analíticos para determinação de picoxistrobina e PIRA por cromatografia eletrocinética capilar micelar e de enrofloxacina por fosforimetria em temperatura ambiente. PUC/Rio, 2008, (Tese de Doutorado).

SPIVAK, S.A.; J. CAMPBELL. Systematic study of steric and spatial contributions to molecular recognition by non-covalent imprinted polymers, **Analyst.** V.126, p. 793, 2001.

SUNDRAVADANA, S., D. Alice, R. SAMIYAPPAN S. KUTTALAM. Determination of Azoxystrobin Residue by UV Detection High Performance Liquid Chromatography in Mango. **J. Braz. Chem. Soc.**, V. 19, N°. 1, p. 60-63, 2008.

SUSHA, A. S. A. M. J., W. J. PARAK, A. L. ROGACH. Luminescent CdTe nanocrystals as ion probes and pH sensors in aqueous solutions. **Colloids and Surfaces A: Physicochem. Eng. Aspects.** V. 281 p. 40–43, 2006.

TARLEY, C.R.T.; SOTOMAR, M.D.P.T.; KUBOTA, L.T. Polímeros biomiméticos em química analítica. Parte 1: Preparo e aplicações de MIP (“Molecularly Imprinted Polymers”) em técnicas de extração e separação. **Quím. Nova**, v.28, 6, 1076-1086, 2005.

VINAS, P., N. CAMPILLO, N. MARTÍNEZ-CASTLHO, M. HERNÁNDEZ-CÓDOBA. Method development and validation for strobilurin fungicides in baby foods by solid-phase microextraction gas chromatography–mass spectrometry. **J. Chromatogr A**. V. 1216, p. 140–146, 2009.

WANG Y., J. CAOB, J.W. A, S. ZENG. Simultaneous determination of quercetin, kaempferol and isorhamnetin accumulated human breast cancer cells, by high-performance liquid chromatography. **J. Pharmaceut. Biom. Anal.** V. 39 p. 328–333, 2005.

WANG G, S. HUI; F. M.B.COUTINHO, G. GALLI, E. CHIELLINI. Copolímeros em Bloco Termotrópicos Poliestireno Poli(Metil-1 ,4-Dioxifenileno-4,4'-Dicarbonil-1,10 Dibenzoil-Oxi-Decano): 1. Síntese e Caracterização. **Polímeros: Ciência e Tecnologia – Artigo técnico científico**. OutlDez -1996.

WEI, Z., LLI-NA MU, Y.HUANG, Z. LIU. Low crosslinking imprinted coatings based on liquid crystal for capillary Electrochromatography. **J. Chromatogr A**. V. 1237, p. 115– 121, 2012.

WEISS, R., MOLINELLI, A., JAKUSCH, M., MIZAIKOFF, B. Molecular imprinting and solid phase extraction of flavonoid compounds. **Bioseparation**. V. 10 p. 379–387, 2002.

WHITHICOMBE, M.J., E.M. RODRIGUEZ, P. VILLAR, E.N. VULSFSON. A new method for the introduction of recognition site functionality into polymers prepared by molecular imprinting - synthesis and characterization of polymeric receptors for cholesterol, **J. Am. Chem. Soc.** V.117, p. 7105, 1995.

WULFF, G., A. SARHAN. The use of polymers with enzyme-analogous structures for the resolution of recemates, **Angew. Chem.** V. 11, p. 341, 1972.

WULFF, G., J HAARER. Enzyme-analogue built polymers. The preparation of defined chiral cavities for the racemic resolution of free sugars, **akromol. Chem.** V.192 p. 1329, 1991.

XIA, Y.Q., T. GUO, M. SONG, B. ZHANG, B. ZHANG. Selective separation of quercetin by molecular imprinting using chitosan beads as functional matrix. **Reactive & Functional Polymers.** V. 66 p.1734–1740, 2006.

XIE, J. L. ZHU, H. LU; L. ZHOU, C. LI, X. J. XU. Direct extraction of specific pharmacophoric flavonoids from gingko leaves using a molecularly imprinted polymer for quercetin. **J. Chromatogr. A.** V. 934 p.1–11. 2001.

XU, J., H. ZHANG, G. CHEN. Carbon nanotube/polystyrene composite electrode for microchip electrophoretic determination of rutin and quercetin in Flos Sophorae Immaturus. **Talanta.** V. 73, p. 932-937, 2007.

YE, L. K. M. The technique of molecular imprinting – principle, state of the art, and future aspects, **J. Inclusion Phenom. Macrocyclic Chem.** V.41, p. 107, 2001.

YEH, W., H., KUO-CHUAN. Amperometric morphine sensing using a molecularly imprinted polymer-modified electrode. **Anal. Chim. Acta.** 542, 76-82, 2005.

YILMAZ, E.; R., O., MOLLER, P., SANCHEZ, D., MOSBACH, K. **J. Mater. Chem.** V. 12, p.1577, 2002.

YIN, J., G. Y., Y.CHEN. Rapid and efficient chiral separation of nateglinide and its L-enantiomer on monolithic molecularly imprinted polymers, **J. Chromatogr. A,** v.1090 p. 68, 2005.

YIN, X., Q. LIU, Y. JIANG, Y. LUO. Development of andrographolide molecularly imprinted polymer for solid-phase extraction. **Spectrochim. Acta Part A.** V. 79, p. 191-196, 2011.

YU, C., HE, YOU-ZHAO, FU, GUO-NI, XIE, H., G., WU-ER. Determination of kanamycin A, amikacin and tobramycin residues in milk by capillary zone electrophoresis with post-column derivatization and laser-induced fluorescence detection. **J. chromatogr. B.** V.877, n<sup>o</sup> 3, p. 333-338, 2009.

ZHANG, X., L. CHEN, Y. XU, H. WANG, Q. ZENG, Q. ZHAO, N. REN, L. DING. Determination of Beta-lactam antibiotics in milk based on magnetic molecularly imprinted polymer extraction coupled with liquid chromatography–tandem mass spectrometry. **J. Chromatogr. B.** V. 878, p. 3421–3426, 2010.

ZHAO, C., T. ZHAO, X. LIU, H. ZHANG. A novel molecularly imprinted polymer for simultaneous extraction and determination of sudan dyes by on-line solid phase extraction and high performance liquid chromatography. **J. Chromatogr. A.** V.1217, p. 6995–7002, 2010.

ZHAO, C., X. GUAN, X. LIA, H. ZHANG. Synthesis of molecularly imprinted polymer using attapulgite as matrix by ultrasonic irradiation for simultaneous on-line solid phase extraction and high performance liquid chromatography determination of four estrogens. **J. Chromatogr.** V.1229, p. 72– 78, 2012.

ZHAO, C., G. JIN, Li-LI CHEN, YI LI, BO YU. Preparation of molecular imprinted film based on chitosan/nafion/nano-silver/polyquercetin for clenbuterol sensing. **Food Chemistry.** V. 129 p. 595–600, 2011.

ZHU, G., J., FAN, Y., GAO, X., GAOA, J., WANG. Synthesis of surface molecularly imprinted polymer and the selective solid phase extraction of imidazole from its structural analogs. **Talanta.** V. 84, p. 1124–1132, 2011.

ZURUTA, A. S. BAYOUDH, P.A.G. CORMACK, L., DAMBIES, J., DEERE, R. BISCHOFF, D.C. SHERRINGTON. Molecularly imprinted solid-phase extraction of cocaine metabolites from aqueous samples, **Anal. Chim.** V. 542 p. 14, 2005.