

Referências bibliográficas

ABURUZ, S.; MILLERSHIP, J.; HEANEY, L.; McELNAY, J. Simple liquid chromatography method for rapid simultaneous determination of prednisolone and cortisol in plasma and urine using hydrophilic lipophilic balanced solid phase extraction cartridges. *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences*. v. 798, n. 2, p. 193-201, 2003.

AMENDOLA, L.; GARRIBA, F.; BOTRÈ, F. Determination of endogenous and synthetic glucocorticoids in human urine by gas chromatography-mass spectrometry following microwavw-assisted derivatization. *Analytica Chimica Acta*, v. 489, p.233-243, 2003.

BOOTH, N.H.; MCDONALD, L.E. *Farmacologia e Terapêutica em Veterinária*. 6^a ed. Rio de Janeiro: Guanabara, 1992.

BRASIL. Ministério da Saúde, Normas e manuais técnicos. Boas práticas de fabricação de produtos farmacêuticos Brasília Centro de Documentação do Ministério da Saúde, 26p., 1983.

BRÄUTIGAM, L.; VETTER, G.; TEGEDER, I.; HEINKELE, G.; GEISSLINGER, G. - Determination of celecoxib in human plasma and rat microdialysis samplas by liquid chromatography tandem mass spectrometry. *Journal of chromatography B*, v. 761, p.203-212, 2001.

BROWER, J.F. Liquid-chromatographic determination of prednisolone in tablets and bulk drugs-interlaboratory study. *Journal of Association of Official Analytical Chemists*. v. 67, n. 4, p. 674-676, 1984.

BYRN, S.R.; XU, W.; NEWMAN, A.W., *Adv. Drug. Deliver.* v. 48, 115. Rev. 2001

CALATAYUD, J. M.; BENITO, C.G. Photochemical derivatization and fluorimetric determination of reserpine in a flow-injection assembly. *Analytica Chimica Acta*, v.245, p. 101-107, 1991.

CALOKERINOS, A.C.; KOUKLI I. I. Continuous-flow chemiluminescence determination of some corticosteroids. *Analyst*, v. 115, n.12, p.1553-1557, 1990.

CARDOSO, C.E. Desenvolvimento e estudo comparativo de metodologias espectrofluorimétrica e voltamétrica para a determinação de talidomida em matrizes diversas, Dissertação de mestrado. Departamento de Química Analítica – PUC-Rio. 2002.

CARDOSO, C.E.; MARTINS, R.O.R.; TELES, C.A.S.; AUCÉLIO, R.Q. Sequential determination of Hydrocortisone and Epinephrine in pharmaceutical formulations via photochemically enhanced fluorescence. *Microchimica Acta*, v. 146, p.79-84, 2004.

CAVICCHIOLI, A.; GUTZ, I.G.R. O uso de radiação ultravioleta para o pré-tratamento de amostras em análise inorgânica. *Química Nova*, v.26, n.6, p. 913-921, 2003.

CHASIN, A.A.M.; CHASIN, M.; SALVADORI, M.R. Validação de métodos cromatográficos em análises toxicológicas. *Revista Farmacêutica e Bioquímica*, Univ. São Paulo, v.30, n.2, p.49-53, 1994.

CHASIN, A.M.; NASCIMENTO, E.S.; NETO, L.M.R.; SIQUEIRA, M.E.P.B.; ANDRAUS, M.R.; SALVADORI, M.C.; FERNÍCOLA, N.A.G.; GORNI, R.; SALCEDO, S. Validação de métodos em análises toxicológicas: uma abordagem geral. *Revista Brasileira de Toxicologia*, v.11, n.1, p.1-6, 1998.

CHENG, X.M. To market, to market – 1994, to market. *Annual reports in medicinal chemistry*, 30:295-317, 1995.

CHENG, X.M. To market, to market – 1995, to market. *Annual reports in medicinal chemistry*, 31:337-355, 1996.

CHO, C.Y.; SHIN, B.S.; YOO, S.D. Sensitive analysis of prednisolone and prednisone in human plasma by reverse phase high-performance liquid chromatography with ultraviolet detection. *Analytical Letters*. v. 36, n. 8, p. 1573-1585, 2003.

CITAC/EURACHEM: Guide to Quality in Analytical Chemistry: An Aid to Accreditation. Citac/ Eurachem, 2002.

DAVIS, L.E. Farmacologia veterinária: Introdução à disciplina. In: BOOTH, N.H.; MCDONALD, L.E. *Farmacologia e Terapêutica em Veterinária*. 6.ed. Rio de Janeiro: Guanabara Koogan, cap.1, p. 2- 5, 1992.

DICIONÁRIO DE ESPECIALIDADES FARMACÊUTICAS 32^a edição – 2003/04, Rio de Janeiro: Ed Publicações científicas.

DOPPENSCHMITT, S.A.; SCHEIDEL, B.; HARRISON, F.; SURMANN, J.P. Simultaneous determination of prednisolone, prednisone acetate and hydrocortisone in human serum by high-performance liquid chromatography. *Journal of Chromatography B-Biomedical applications*. v. 674, n. 2, p. 237-246, 1995.

EL-SAHARTY, Y.S.; HASSAN, N.Y.; METWALLY, F.H. Simultaneous determination of terbinafine HCL and triamcinolone acetone by UV derivative spectrophotometry and spectrodensitometry. *Journal of Pharmaceutical and Biomedical Analysis*. v. 28, p.569-580, 2002.

EURACHEM: The Fitness for Purpose of Analytical Methods: A Laboratory Guide to Method Validation and Related Topics. 1th.ed. Teddington, 1998. ISBN 0-948926-12-0.

FEDERAÇÃO EQUESTRE INTERNACIONAL: Controle de Medicação: Análises para detecção de substâncias proibidas em competições. Disponível em:<<http://www.equisport.pt/45-49/art482.htm>> Acesso em 27/01/2003.

FRERICHS, V.A.; TORNATORE, K.M. Determination of the glucocorticoids prednisone, prednisolone, dexamethasone, and cortisol in human serum using liquid chromatography coupled to tandem mass spectrometry. *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences*. v. 802, n. 2, p. 329-338, 2004.

GAILLARD, Y.; VAYSETTE, F.; PÉPIN, G. Compared interest between hair analysis and urinalysis in doping controls . Results for amphetamines, corticosteroids and anabolic steroids in racing cyclists. *Forense Science International*. v. 107, p. 361-379, 2000.

GALATISIS, P. To market, to market – 1996, to market. *Annual reports in medicinal chemistry*, 32:305-3327, 1997.

GALATISIS, P. To market, to market – 1997, to market. *Annual reports in medicinal chemistry*, 33:327-353, 1998.

GALLEGO, J.M.L; ARROYO, J.P. Determination of prednisolone acetate, sulfacetamide and phenylefrine in local pharmaceutical preparations by micellar electrokinetic capillary chromatography. *Journal of Pharmaceutical and Biomedical Analysis*. v. 31, n. 5, p. 873-884, 2003.

GALLEGO, J.M.L.; ARROYO, J.P. Determination of prednisolone and the most important associated compounds in ocular and cutaneous pharmaceutical preparations by micellar electrokinetic capillary chromatography. *Journal of Chromatography B-Analytical Technologies in the Biomedical and Life Sciences*. v. 784, n. 1, p. 39-47, 2003.

GALLEGO, J.M.L.; ARROYO, J.P. Determination of prednisolone, naphazoline, and phenylephrine in local pharmaceutical preparations by micellar electrokinetic capillary chromatography. *Journal of Separation Science*. v. 26, n. 9-10, p. 947-952, 2003.

GATTI, R.; GIOIA, M.G.; DI PIETRA, A.M.; CAVRINE, V. HPLC-fluorescence determination of unconjugated estrogens in pharmaceuticals. *Journal of Pharmaceutical and Biomedical Analysis*, Amsterdam, v. 18, p. 187-192, 1998.

GENÉRICO INFO. Disponível em: <<http://www.genericoinfo.hpg.ig.com.br/noticias.htm>> Acesso em 02/09/04.

GOODMAN, L.S.; GILMAN, A.G. *As bases farmacológicas da terapêutica*. 7^a ed. Rio de Janeiro: Editora Guanabara, 1987.

GORDON, G.; WOOD, P.R. Determination of triamcinolone acetonido in cream and suspension formulations by High-Performance Liquid-Chromatography. *Analyst*. v. 101, n. 1208, p. 876-882, 1976.

GÖRÖG, S. Recent advances in the analysis of steroid hormones and related drugs. *Analytical Sciences*, v. 20, p.767-782, 2004.

GOTZMANN, A.; BREDEHÖFT, M.; THEVIS, M.; SCHANZER, W. Detection of Prednisolone, Prednisone and 20 β -Dihydroprednisolone in equine urine by means of LC/MS/MS. In: INTERNACIONAL CONFERENCE OF RACING ANALYSTS AND VETERINARIANS, 14, 2002, Florida. Proceedings. Florida, p.176-179, 2003.

HANDBOOK OF CHEMISTRY AND PHYSICS. 75^a ed. USA:CRQ Press, p. 15 e 43, 1994

HARPER, H.A.; RODWELL, V.W.; MAYES, P.A. *Manual de Química Fisiológica*. 5^a ed. São Paulo: ATHENEU, 1982.

HARVEY, R.A.; CHAMPE, P.C.; MYCEK, M.J. *Farmacologia Ilustrada*. 2^a ed. Porto Alegre: ARTMED, 1998.

HEWLETT-PACKARD. HP 1046A Fluorescence Detector: user's guide. Germany, 1994.

IGLESIAS, Y.; FENTE, C.; VAZQUEZ, B.I.; FRANCO, C.; CEPEDA, A.; MAYO, S. Application of the luminol chemiluminescence reaction for the determination of nine corticosteroids. *Analytica Chimica Acta*, v. 468, n.1, p.43-52, 2002.

IGLESIAS, Y.; FENTE, C.; VAZQUEZ, B.I.; FRANCO, C.; CEPEDA, A.; MAYO, S. Chemiluminescence detection of nine corticosteroids in liver. *Analyst*, v. 125, n.11, p.2071-2074, 2000.

INGLE, J.D., CROUCH, S.R. *Spectrochemical analysis*. 1^a ed. New Jersey: Prentice Hall, 1988.

JABLONSKY, A. *Molecular Luminescence Spectroscopy*. Z Phhys. 1935

KATAYAMA, M.; MASUDA, Y.; TANIGUCHI, H. Determination of corticosteroids in plasma by high-performance chromatography after precolumn derivatization with 2-(4-carboxyphenyl)-5-6-dimethylbenzimidazole. *Journal of chromatography-biomedical applications*, v. 612, n.1, p.33-39, 1993.

KATZUNG, B.G. *Farmacologia básica & clinica*. 5^a ed. Rio de Janeiro: Editora Guanabara, 1994.

KEDORHACKMANN, E.R.M.; GIANOTTO, E.A.S.; SANTORO M.I.R.M. Determination of triamcinolone acetonido and salicylic acid in pharmaceutical formulations by High Performance Liquid Chromatography. *Pharmazie*. v. 51, n. 1, p. 63-63, 1996.

KOUPAI-ABYAZANI, M. R.; YU, N.; ESAW, B.; LAVIOLETTE, B. Determination of triancinolona acetonido in equine serum and urine by liquid chromatografy-atmospheric pressure ionisation mass spectrometry. In: *INTERNACIONAL CONFERENCE OF RACING ANALYSTS AND VETERINARIANS*, 10, 1994, Stockholm. *Proceedings*. Suffolk, R&W, 1995. p. 209-213.

LONG, G.L.; WINEFORDNER, J.D. Limit of detection: A closer look at the IUPAC definition. *Analytica Chimica Acta*, v. 55, p.712A-724A, 1983.

MANUAL MERCK ON-LINE. Disponível em: <http://www.msd-brazil.com/msd43/m_manual> Acesso em 12/06/04.

MASON, S.R.; WARD, L.C.; REILLY, P.E.B. Fluorimetric detection of serum corticosterone using high-performance liquid chromatography. *Journal of Chromatography*, v. 581, p. 267-271, 1992.

MATTINGLY, D. A simple fluorimetric method for the estimation of free 11-hydroxycorticoids in human plasma. *Journal Clin.Path.*, v. 15, p.374-379, 1962.

MATYSOVA, L.; HAJKOVA, R.; SICHA, J.; SOLICH, P. Determination of methylparaben, propylparaben, triamcinolone acetone and its degradation product in a topical cream by RP-HPLC. *Analytical and Bioanalytical Chemistry*. v. 376, n. 4, p. 440-443, 2003.

NETO, L.M.R.; SALVADORI, M.C.; SPINOSA, H.S. Immunoaffinity chromatography in the detection of Dexamethasone in equine urine. *Journal of Chromatography Science*, v. 35, p. 543-551, 1997.

NETO, L.M.R.; SALVADORI, M.C.; SPINOSA, H.S. Determinação de hidrocortisona em urina de cavalo de corrida empregando cromatografia a líquido de alta eficiência com detetor de arranjo de diodos. *Revista Brasileira de Toxicologia*, v.12, n.2, p. 100-104, 1999.

NETO, L.M.R.; SALVADORI, M.C.; SPINOSA, H.S. Panorama atual da análise de corticosteróides no controle antidopagem de cavalos de corrida. *Revista Brasileira de Toxicologia*, v.9, n.2, p. 13-17, 1996.

NETO, L.M.R.; SALVADORI, M.C.; SPINOSA, H.S.; LOPEZ, N.M.R. The use of ELISA tests and immunoaffinity chromatography combined with reversed-phase high-performance liquid chromatography for Dexamethasone detection in equine urine. *Journal of Analytical Toxicology*, v.21, p. 393-396, 1997.

NEUFELD, E.; CHAYEN, R.; STERN, N. Fluorescence derivatisation of urinary corticosteroides for high-performance liquid chromatography analysis. *Journal of Chromatography B*, v. 718, p. 273-277, 1998.

NOZAKI, O. Steroid analysis for medical diagnosis. *Journal of Chromatography A*, v. 935, p. 267-278, 2001.

PAN, Z.T.; YU, J.P. Fluorescent spectrometry of triamcinolone acetone. *Chinese Journal of analytical chemistry*, v. 32, n.3, p.338-341, 2004.

PAZZAGLI, M.; MESSERI, G.; SALERNO, R.; CALDINI, A.L.; TOMMASI, A.; MAGINI, A.; SERIO, M. Luminescent immunoassay (LIA) methods for steroid hormones. *Talanta*, v. 31, n. 10B, p.901-907, 1984.

PEREIRA, H.M.G.; MARQUES, M.A.S.; CARDOSO, J.N.; NETO, F.R.A. Análise de glicocorticosteróides por CG-EM: Uma nova abordagem de derivação para o controle de dopagem no esporte. *Química Nova*, v. 25, n. 6B, p.1096-1102, 2002.

POPOT, M. A.; GARCIA, P. Detection of Triamcinolone, Betamethasone and Methylprednisolone in the horse: effect on urine cortisol and cortisol metabolite concentrations. In: INTERNACIONAL CONFERENCE OF RACING ANALYSTS AND VETERINARIANS, 14, 2002, Florida. Proceedings. Florida, 2003. p.408-412.

PORTUGAL, H.; PAULI, A.M.; PASTOR, J. Spectrofluorimetric determination of prednisolone and prednisone. *Analisis*, v. 17, n.5, p.275-279, 1989.

PRISTA, L.N. Segurança e eficácia medicamentosa e controle de qualidade. *Revista Portuguesa de Farmácia*. v.38, n.3, p.1-8, 1988.

REIS, A.M.M.; GOMES, M.J.V.M. Ciências Farmacêuticas – Uma abordagem em farmácia hospitalar. São Paulo: ATHENEU, [19--].

REIS, A.M.M.; RESENDE, D.C.; CAMPOS, L.M. Testes de controle de qualidade de medicamentos. Belo Horizonte: SBRAFH, 22p, 1999.

RODCHENKOV, G.M.; URALETS, V.P.; SEMENOV, V.A. Gas-chromatographic and mass-spectrometric determination of triamcinolone and its metabolites in human biological-fluids. *Khimiko-farmatsevticheskii Zhurnal*. v. 22, n. 5, p. 622-625, 1988.

RODCHENKOV, G.M.; URALETS, V.P.; SEMENOV, V.A.; GUREVICH, V.A. Gas chromatographic and mass spectral study of betamethasone synthetic corticosteroid metabolism. *Journal of Chromatography*. v. 432, p. 283-289, 1988.

RODCHENKOV, G.M.; URALETS, V.P.; SEMENOV, V.A.; VEDENIN, V.P. Characterization of prednisone, prednisolone and their metabolites by gas chromatographic-mass spectrometry. *Journal of Chromatography*. v. 565, p. 45-51, 1991.

ROSENBERG F.J.; SILVA, A.B.M. As farmacopéias e o laboratório oficial de controle de qualidade. In: Bonfim JBA, Mercucci VB eds. A construção da política de medicamentos. São Paulo: Hucitec-Sobravime, p. 251-255, 1997.

SAHA, R.N.; SAJEEV, C.; JADHAV, P.R.; PATIL, S.P.; SRINIVASAN, N. Determination of celecoxib in pharmaceutical formulations using UV spectrophotometry and liquid chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, v. 28, p. 741-751, 2002.

SALVADORI, M. C. Aspectos gerais do doping no cavalo de esporte. *Cad.Téc.Esc.Vet., UFMG*, n.19, p.77- 86, 1997.

SCHÖNBERGER, F.; HEINKELE, G.; MÜRDTER, T.E.; BRENNER, S.; KLOTZ, U.; HOFMANN, U. Simple and sensitive method for the determination of

celecoxib in human serum by high-performance liquid chromatography with fluorescence detection. *Journal of chromatography B*, v. 768, p. 255-260, 2002.

SCHULMAN, S.G. *Fluorescence and phosphorescence spectroscopy: physicochemical principles and practice*. 1^a ed. New York: Pergamon Press, 1977.

SCHULMAN, S.G. *Molecular Luminescence Spectroscopy - Methods and Applications: Part 1 (Chemical analysis v.77)*. New York: A WILEY-INTERSCIENCE PUBLICATION, 1985.

SHIBATA, N.; HAYAKAWA, T.; TAKADA, K.; HOSHINO, N.; MINOUCHI, T.; YAMAJI, A. Simultaneous determination of glucocorticoids in plasma or urine by high-performance liquid chromatography with precolumn fluorimetric derivatization by 9-anthroyl nitrile. *Journal of chromatography B-analytical technologies in the biomedical and life sciences*, v. 706, n.2, p.191-199, 1998.

SPINELLI, E.; SILVA, O.A. Imunoensaio de fluorescência polarizada e sua aplicação na detecção de usuários de cannabis. *Revista Brasileira de Toxicologia*, v.7, n.1/2, p. 37-43, 1994.

STANLEY, S. M. R.; RODGERS, J. P. Use of immunoaffinity chromatography with HPLC-particle beam/MS and CG/MS for the confirmatory analysis of corticosteroids in equine urine. In: *INTERNACIONAL CONFERENCE OF RACING ANALYSTS AND VETERINARIANS*, 10, 1994, Stockholm. Proceedings. Suffolk, R&W, 1995. p. 226-232.

TAYLOR, B.N. *Guide for the use of the International System of Units (SI)*. NIST Special Publication, 1995.

TEALE, P.; HINES, S.; DUMASIA, M.; MARSHALL, D.; HOUGHTON, E. Confirmation of metabolite of Triamcinolone Acetonido: a comparison of LC/MS and GC/MS data. In: *INTERNACIONAL CONFERENCE OF RACING ANALYSTS AND VETERINARIANS*, 14, 2002, Florida. Proceedings. Florida, 2003. p.435-441.

THE MERCK INDEX: *Enciclopédia of Chemicals, Drugs and Biologicals*. 11^a ed.

THE PHARMACEUTICAL SOCIETY OF GREAT BRITAIN. *Clarke's Isolation and Identification of Drugs in pharmaceuticals, body fluids and post-mortem material*. 2th.ed.London, ISBN 0-85369-166-5, p. 474, 475, 917, 918, 1036 e 1037, 1986.

THE UNITED STATES PHARMACOPEIA THE NATIONAL FORMULARY – USP XXII, NF XVII, p.1395-1396 e 1128-1130, 1990.

TOBIN, T. Drugs and the performance horse. Illinois: Charles C. Thomas, 1981.

TOBIN, T.; SANS, R.A. Doping em cavalos, 1999. Disponível em: <<http://www.bichoonline.com.br/artigos>> Acesso em 27/01/2003.

UDENFRIEND, S. Fluorescence Assay in Biology and Medicine. V.2, New York: Academic Press, 1969.

WEHRY, E.L. Modern fluorescence spectroscopy. 2 New York: Penum Press, 1976.

XU, Y.M.; WONG, G.Y. Simultaneous determination of lignocaine hydrochloride, chlorhexidine gluconate, and triamcinolone acetone in suspension by reversed-phase HPLC. Journal of Liquid Chromatography & Related Technologies. v. 22, n. 13, p. 2071-2091, 1999.

YAMAGUCHI, M.; ISHIDA, J.; YOSHITAKE, T.; NAKAMURA, M. Determination of prednisolone and prednisone in plasma by liquid chromatography with fluorescence detection. Analytica Chimica Acta, v. 242, p.113-116, 1991.

YAP, B.K.; JOHNSTON, G.A.R.; KAZLAUSKAS, R. Routine screening and quantification of urinary corticosteroids using bench-top gas chromatography-mass-selective detection. Journal of Chromatography, v. 573, p. 183-190, 1992.

YOSHITAKE, T.; HARA, S.; YAMAGUCHI, M.; NAKAMURA, M.; OHKURA, Y.; GOROG, S. Measurement of 21-hydroxycorticosteroids in human and rat sera by high-performance liquid chromatography with fluorimetric detection. Journal of Chromatography, v. 489, p. 364-370, 1989.

ZANINI, A. C.; OGA, S. Farmacologia Aplicada. 5^a ed. São Paulo: Atheneu, 1994.

Glossário

Analgesia – perda de sensibilidade.

Catabolização – processo metabólico destrutivo, por meio do qual células do organismo transformam substâncias complexas em outras mais simples.

Hiperalgia – exagero da sensibilidade à dor; hiperalgia.

Leucócitos – células sanguíneas incolores envolvidas na defesa orgânica celular e imunitária; glóbulos brancos.

Nefrotoxicidade – toxicidade para os rins.

Quimiotático – vem de quimiotaxia que quer dizer ação atrativa ou repulsiva demonstrada por certas células vivas em relação a outras células ou substâncias que exercem sobre aquelas uma influência química; quimiotactismo.

Termistor – resistor cuja resistência decresce fortemente com a elevação de temperatura, e que é, em regra, constituído por um semicondutor.